

When the objective is prevention of iron deficiency in pregnant women, an amount of 15mg to 30mg iron per day are adequate to meet the requirement in 2nd & 3rd trimester"

- Harrison's, 15th Ed. Vol.1.

^RHEMGARD-XT SYRUP

Usual dose: 1-2 tsf daily

%HEMGARD-XT TABLET

Usual dose: 1 Tablet OD

B12, Folic acid, Iron is the need in Therapy of Oral Iron

HEMGARD-XT

Each Tablet contains Ferrous Ascorbate Eqv. to elemental Iron 100mg + Folic Acid 1.5mg Each 5ml Syrup contains Ferrous Ascorbate Eqv. to elemental Iron 30mg + Folic Acid 0.55mg + Vitamin B12 7.5mcg

Ferrous Ascorbate

- Better iron absorption
- No constipation / G.I. complications
- Assured hemoglobin response

"Combination of Ferrous compound with reducing sugars or ascorbate prevents or returds oxidation of Ferrous molecule resulting an increased absorption of iron. Ascorbate prevents oxidation of ferrous iron, and Ferrous ascorbate release ferrous ions directly to gastric mucosa from ascorbate complex."

 Dept. Of Pharmaceutical Analysis & Analytical Chemistry. University of Leiden, Netherlands



HEM